



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Treatment Algorithms in Stage IV Melanoma

Espinosa, Enrique ; Grob, Jean-Jacques ; Dummer, Reinhard ; Rutkowski, Piotr ; Robert, Caroline ; Gogas, Helen ; Kefford, Richard ; Eggermont, Alexander M M ; Martin Algarra, Salvador ; Hauschild, Axel ; Schadendorf, Dirk

Abstract: The molecular classification of melanoma and the advent of new drugs are changing the paradigm of therapy for advanced melanoma. A review of the recent key studies was performed, followed by a discussion in an expert forum. The aim of this review was to generate a therapeutic algorithm for stage IV melanoma. Tumor genotyping for BRAF and/or KIT should be performed before selection of therapy. For most BRAF-mutated melanoma patients and particularly those with a high tumor load, vemurafenib or other BRAF inhibitors such as dabrafenib are the treatment of choice. KIT inhibitors can be effective in KIT-mutant tumors, especially in those patients with mutations at exons 11 and 13. Ipilimumab is a good option for patients with nontargetable or nondetected mutations and those who progress under therapy with vemurafenib or a KIT inhibitor. There is still a role for conventional chemotherapy either as first-line treatment in BRAF wild-type patients or as salvage therapy in second or third line, or after other treatment modalities. Participation in clinical trials is strongly encouraged, either in first or in subsequent lines. New therapeutic options for advanced melanoma are guided by tumor genotyping. The current therapeutic algorithm includes kinase inhibitors, anti-CTLA4 therapy, immunotherapy, and chemotherapy, depending on the tumor genotype and response to previous treatments. Participation in clinical trials should always be encouraged because the treatment goal is long-term survival and potential cure in a subset of patients.

DOI: <https://doi.org/10.1097/MJT.0b013e31829e885c>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-93243>

Journal Article

Published Version

Originally published at:

Espinosa, Enrique; Grob, Jean-Jacques; Dummer, Reinhard; Rutkowski, Piotr; Robert, Caroline; Gogas, Helen; Kefford, Richard; Eggermont, Alexander M M; Martin Algarra, Salvador; Hauschild, Axel; Schadendorf, Dirk (2015). Treatment Algorithms in Stage IV Melanoma. *American Journal of Therapeutics*, 22(1):61-67.

DOI: <https://doi.org/10.1097/MJT.0b013e31829e885c>

Treatment Algorithms in Stage IV Melanoma

Enrique Espinosa, MD,^{1*} Jean-Jacques Grob, MD,² Reinhard Dummer, MD,³ Piotr Rutkowski, MD, PhD,⁴ Caroline Robert, MD, PhD,⁵ Helen Gogas, MD,⁶ Richard Kefford, MD,⁷ Alexander M. M. Eggermont, MD, PhD,⁵ Salvador Martin Algarra, MD, PhD,⁸ Axel Hauschild, MD,⁹ and Dirk Schadendorf, MD¹⁰

The molecular classification of melanoma and the advent of new drugs are changing the paradigm of therapy for advanced melanoma. A review of the recent key studies was performed, followed by a discussion in an expert forum. The aim of this review was to generate a therapeutic algorithm for stage IV melanoma. Tumor genotyping for *BRAF* and/or *KIT* should be performed before selection of therapy. For most *BRAF*-mutated melanoma patients and particularly those with a high tumor load, vemurafenib or other *BRAF* inhibitors such as dabrafenib are the treatment of choice. *KIT* inhibitors can be effective in *KIT*-mutant tumors, especially in those patients with mutations at exons 11 and 13. Ipilimumab is a good option for patients with nontargetable or nondetected mutations and those who progress under therapy with vemurafenib or a *KIT* inhibitor. There is still a role for conventional chemotherapy either as first-line treatment in *BRAF* wild-type patients or as salvage therapy in second or third line, or after other treatment modalities. Participation in clinical trials is strongly encouraged, either in first or in subsequent lines. New therapeutic options for advanced melanoma are guided by tumor genotyping. The current therapeutic algorithm includes kinase inhibitors, anti-CTLA4 therapy, immunotherapy, and chemotherapy, depending on the tumor genotype and response to previous treatments. Participation in clinical trials should always be encouraged because the treatment goal is long-term survival and potential cure in a subset of patients.

Keywords: melanoma, advanced disease, *BRAF* inhibitor, MEK inhibitor, anti-CTLA4

INTRODUCTION

The management of metastatic melanoma has yielded disappointing results until recent times. Some patients obtained a benefit from regional treatments, but the majority of them required systemic therapy that was largely ineffective. Options included immunotherapy in selected centers (and for selected patients), clinical trials with new drugs, and, most commonly, chemotherapy.

Both new knowledge of the molecular biology of melanoma and the advent of ipilimumab and specific kinase inhibitors have dramatically changed this landscape. Melanoma is no longer envisioned as one single tumor but a constellation of diseases having specific molecular features. This new understanding allowed the development of *BRAF* inhibitors, for instance, and will likely foster the identification of other effective and specific drugs. As new compounds enter clinical

¹Service of Oncology, Hospital La Paz, Madrid, Spain; ²Department of Dermatology, Hôpital Ste Marguerite, Marseille, France; ³Department of Dermatology, University Hospital of Zurich, Zurich, Switzerland; ⁴Department of Sarcoma and Melanoma, M. Skłodowska-Curie Memorial Cancer Center and Institute of Oncology, Warsaw, Poland; ⁵Department of Dermatology (CR), and General Director (AE), Institut Gustave Roussy, Villejuif Cedex, France; ⁶First Department of Medicine, University of Athens Medical School, Athens, Greece; ⁷Department of Oncology, Westmead Hospital and Melanoma Institute Australia, University of Sydney, Sydney, Australia; ⁸Service of Oncology, Clínica Universitaria de Navarra, Pamplona, Spain; ⁹Department of Dermatology, University of Kiel, Kiel, Germany; and ¹⁰Department of Dermatology, University Hospital Essen, Essen, Germany.

The authors have no conflicts of interest to declare.

*Address for correspondence: Service of Oncology, Hospital La Paz, Paseo de la Castellana, 261, Madrid 28046, Spain. E-mail: eespinosa00@hotmail.com

trials and eventually become widely available, decision trees will be required to use them correctly.

This review summarizes evidence that may help in making clinical decisions and proposes an algorithm to use systemic treatments. A review of the recent key studies was performed, followed by a discussion in an expert forum. Local therapies will continue to play an important role in the management of regional disease but will not be covered in this review.

OPTIONS AND OUTCOME BEFORE THE ERA OF NEW DRUGS

The prognosis of patients with stage IV melanoma has remained unchanged for decades. A retrospective analysis published in 1983 showed a 1-year survival rate of 40% for stage IVA and 11% for stage IVC disease.¹ In 2000, the Eastern Cooperative Oncology Group reported a median overall survival of 8–10 months in patients with soft tissue or lung metastases and 6 months in those with visceral dissemination.² Numerous efforts to improve this outcome with the use of combined chemotherapy and immunotherapy proved to be unsuccessful. A systematic review of 41 randomized clinical trials revealed that combination regimens produced higher response rates, but at the cost of increased toxicity and with no benefit in overall survival.³ These trials had used single-agent chemotherapy, combination chemotherapy, interleukin-2 with or without interferon or combinations of chemotherapy, and immunotherapy. In view of the limited success of complex schemes, single-agent therapy with dacarbazine, which achieves responses in 7%–15% of patients, was accepted as a reasonable standard of care in many institutions. Temozolomide and fotemustine were also used because they had similar activity and a favorable toxicity profile, and by crossing the blood-brain barrier, they have activity in brain-metastasized patients as well.^{4–6}

Some types of immunotherapy stand out as exceptions to these poor results. Interleukin-2 yields complete remissions in 6% of patients, most of which are long-term survivors.⁷ Because of the side effects in selected patients associated with this drug, it is usually restricted to young patients with excellent performance status and no comorbid conditions. Therapy with interleukin-2 is offered in selected centers with experience in the management of toxicity.

Adoptive cell therapy consists of the infusion of the patient's own tumor infiltrating lymphocytes after ex vivo expansion or antigen-specific T lymphocytes, that is, T-cell clones. The infusion is usually preceded by an immune-depleting regimen including chemotherapy or total-body irradiation. The procedure should be restricted

to highly selected patients and requires a technology that is not widely available. In experienced hands, 3-year survival rates of up to 42% have been reported.^{8,9}

ANTI-CTLA4 THERAPY

Ipilimumab is a fully humanized monoclonal antibody directed to CTLA4. Long-term survival in some patients was observed in phase 2 studies, although overall response rates were low.¹⁰ A double-blinded phase 3 study in second line compared ipilimumab (3 mg/kg every 3 weeks) versus ipilimumab plus a gp100 vaccine versus the vaccine alone. Overall survival favored the ipilimumab arms, with 45.6% and 23.5% of the patients remaining alive at 1 and 2 years, respectively.¹¹ Another phase 3 study in first line compared dacarbazine versus dacarbazine plus ipilimumab (10 mg/kg every 3 weeks). Again, survival rates were superior in the ipilimumab arm: 47% versus 36%, 28% versus 18%, and 20% versus 12% at 1, 2, and 3 years, respectively.¹² Median overall survival was 11 months for the ipilimumab plus dacarbazine combination versus 9 months for dacarbazine alone. Although ipilimumab showed little benefit in overall response rate or progression-free survival, a survival plateau appeared after 2 years of follow-up, with 20% of the patients remaining alive in the long term. Based on these results, ipilimumab 3 mg/kg has been approved by the Food and Drug Administration (FDA) for first and second line, whereas the European Medicine Agency (EMA) and the Therapeutic Goods Administration in Australia approved it for second-line therapy. Interestingly, long-term benefit with ipilimumab does not seem to depend on the BRAF status.¹³

Ipilimumab was the first agent to demonstrate a significant overall survival benefit in metastatic melanoma but generated new issues. Standard criteria for the evaluation of response are not accurate for patients treated with ipilimumab, because initial lymphocyte infiltration in the tumor may increase tumor size or lead to the appearance of new lesions in computed tomography images (pseudoprogression). As a consequence, immune-related response criteria have been developed.¹⁴ Furthermore, ipilimumab is associated with a new toxicity profile of immune-related side effects, mostly cutaneous and gastrointestinal, which requiring specific education and training of treating clinicians.

Future studies with the drug may be directed to the identification of predictive factors and the combination of ipilimumab with other drugs, such as BRAF inhibitors. Other agents that enhance the activity of the immune system are under development, such as those targeting PD1 and PD1 ligand. In the multicenter phase 1 trial, anti-PD1 ligand antibody was given

intravenously to a total of 207 patients as of February 2012. These included 55 with melanoma. Nine melanoma patients (of 52 who could be evaluated) showed an objective response, either complete or partial. Responses were durable in many patients, providing evidence that this line of therapy should be part of melanoma treatment in the future.^{15,16}

SPECIFIC THERAPY IN BRAF-MUTATED MELANOMA

Approximately 40% of melanomas present mutations in *BRAF*, usually V600E and less commonly V600K and others.^{17,18} This has allowed the development of specific kinase inhibitors. A diagram showing the role of *BRAF* in melanoma is given in Figure 1.¹⁹

BRAF inhibitors

Vemurafenib is a selective *BRAF* inhibitor that targets the V600 mutant forms of the *BRAF*.^{20,21} Detection of the *BRAF* mutation can be done in paraffin-embedded specimens. The drug is active only in tumors harboring this kind of mutation, which needs to be detected before treatment initiation. Whenever possible, the mutation should be detected in metastatic tissue, as disparities have been reported between primary and metastatic lesions in some cases.²²




FIGURE 1. The BRAF-MEK pathway. Inappropriate activation of growth factor receptors (eg, KIT) or mutations of the *BRAF* gene can lead to constant activation of MEK 1 or 2, with resultant effects on cells: increased proliferation, survival, and propensity for invasion. The diagram is adapted from Meier et al.¹⁹

www.americantherapeutics.com

A phase 2 study of vemurafenib in second/third line showed objective responses in 53% of the patients and a median progression-free survival of 6.7 months.²³ The median overall survival was 15.9 months. The most common grade 3 adverse events requiring dose reduction were arthralgia, rash, UV A-dependent photosensitivity, fatigue, and elevated transaminases in 2%–5% of the patients.^{23,24} Skin lesions such as squamous cell carcinoma and keratoacanthoma developed in 24% of the patients and were managed by surgical excision.

A subsequent phase 3 trial compared vemurafenib—960 mg every 12 hours—to dacarbazine in first line.²⁵ The first interim analysis showed a response rate of 48% for vemurafenib (vs. 5% for dacarbazine) and a median progression-free survival of 5 months (vs. 1.6). Based on these results, the FDA and the EMA have approved vemurafenib for the treatment of *BRAF*-mutated metastatic melanoma.

Vemurafenib is being investigated in a phase 1/2 study in combination with ipilimumab for advanced disease. There are also ongoing studies in patients with poor performance status or with brain metastases.^{26–28}

Other *BRAF* inhibitors are being developed. Of these, dabrafenib (GSK118436) obtained results similar to those of vemurafenib in a phase 1/2 study.²⁹ A recent comparison of dabrafenib and dacarbazine in the BREAK-3 trial showed that dabrafenib demonstrated significant improvement in progression-free survival and overall relapse rate compared with dacarbazine.³⁰ Dabrafenib also showed significant activity in brain metastases with an overall response rate of 39% as first-line treatment and 31% in those who had previous brain radiotherapy.³¹

MEK inhibitors

MEK is a downstream target of the *BRAF* pathway, which is depicted in Figure 1. MEK mutation has been described as a possible mechanism of resistance to *BRAF* inhibitors.³² Patients developing resistance to *BRAF* inhibitors frequently show reactivation of the MAPK pathway,³³ and a specific MEK inhibitor, trametinib (GSK1120212), has shown activity in patients who were progressing with a *BRAF* inhibitor.³⁴ A recent phase 3 study, the METRIC trial, compared trametinib versus chemotherapy with either dacarbazine or paclitaxel in patients with *BRAF*-mutant nonresectable melanomas. The results showed that trametinib conferred significant improvement in progression-free survival and overall survival compared with either chemotherapeutic modality.³⁵

Combinations of specific inhibitors

Considerable effort is being dedicated to study the resistance to *BRAF* inhibitors and a number of mechanisms have been proposed. For instance, preclinical

American Journal of Therapeutics (2014) 0(0)

models indicate the importance of MEK dependency in *BRAF*-mutant melanoma and suggest that a combination of *BRAF* and MEK inhibitors could prevent the emergence of resistance.³⁶ A phase 1/2 study of the *BRAF* inhibitor dabrafenib and the MEK inhibitor trametinib has shown encouraging activity with lower than expected toxicity. Among the 77 melanoma patients treated with this therapy combination, the overall response rate was 56% and the overall PFS was 7.4 months. The authors concluded that the combination had an acceptable safety profile along with antitumor efficacy.³⁷ Phase 3 trials comparing the MEK/*BRAF* inhibitor combination to *BRAF* inhibitor alone are currently underway.

Other mechanisms of resistance, independent of the *BRAF* or MEK pathways, have been described and point to new possible drug combinations.^{38–42} At this time, the best strategy to prevent or overcome resistance to *BRAF* inhibitors remains unknown and will be the aim of future clinical trials. Given the heterogeneity of melanoma, it is unlikely that one particular combination will work in all cases.

KIT inhibitors

Approximately 15% of mucosal (with an especially high mutation rate in vulvovaginal melanomas⁴³) and 23% of acral melanomas have a mutation or amplification in *KIT*,⁴⁴ which could allow therapy with specific inhibitors. A phase 2 trial of imatinib 800 mg/d in Chinese patients reported a partial response rate of 23% and stabilization in 30% of patients.⁴⁵ There is a clear hierarchy of mutations according to the response to targeted drugs with some mutations being completely resistant.⁴⁶ Most imatinib-sensitive mutations in melanoma are found in exons 11 and 13. In a phase 2 study from 28 patients with *KIT* mutation or amplification, 6 responded to the drug.⁴⁶ All of them had tumors with L576P or K642E mutations, the most common *KIT* mutations in melanoma. Experience in Western patients is more limited.⁴⁶ Likewise, the incidence of *KIT* mutation could differ depending on ethnicity. Ongoing studies (NCT01395121) have been launched for another specific inhibitor, nilotinib, in this population.

CONVENTIONAL CHEMOTHERAPY AND OTHER COMPOUNDS

Twenty percent of melanomas harbor mutations in N-RAS, whereas uveal melanomas typically have mutations in *GNAQ* or *GNA11*, and other melanomas yet show a variety of other mutations (PI3K pathway, *MITE*, *CDKs*, etc.).⁴⁷ MEK inhibitors, such as MEK162, could play a role in some of these less common subgroups of

melanoma. This selective inhibitor of MEK1 and MEK2 showed clinical efficacy with good tolerability in an open-label phase 2 trial, among patients bearing *BRAF*- and *NRAS*-mutated melanoma.⁴⁷ Further basic research and multi-institutional cooperation will be critical in the future management of these subgroups.

However, chemotherapy has traditionally been associated with poor overall results, but some patients clearly benefit from this approach. For this reason, chemotherapy will still have a place as rescue strategy in second or third line and in patients with nonmutated melanoma.


A THERAPEUTIC ALGORITHM

As therapy for advanced melanoma becomes more complex, it is important to delineate treatment algorithms. Registration labels and drug availability will drive the sequence. At this time, ipilimumab and vemurafenib have been approved by regulatory agencies, although cost constraints may limit access in some countries. Other new drugs and combination options mentioned above remain experimental.

Considering the high response rate associated with vemurafenib, tumor genotyping is the first logical step in the algorithm (Figure 2). *BRAF* mutation should be searched for in all patients with metastatic melanoma.⁴⁸ An automated polymerase chain reaction-based diagnostic test has been developed and is approved by the FDA.⁴⁹ This test is very sensitive to detect V600E mutations but is less sensitive for non-V600E mutations. For this reason, if a mutation in *BRAF* is not found, ordering a second analysis by an alternative method could be justified. Acral and mucosal melanomas should also be tested for mutations in *BRAF* and *KIT*. *NRAS* mutational status should be determined in wild-type *BRAF* melanoma, as MEK inhibitors are now under clinical evaluation in *NRAS*-mutated disease.⁴⁷

Vemurafenib—and probably dabrafenib—is the preferred option in patients with *BRAF*-mutated melanoma. However, ipilimumab could be considered in low-risk patients, that is, those with low tumor burden and excellent performance status. Some of these patients could become long-term survivors, whereas the remaining would receive vemurafenib on progression with ipilimumab. As ipilimumab requires time to produce an effective response, patients with short life expectancy could be better served with other options.

Imatinib or another *KIT* inhibitor is indicated if a *KIT* mutation is detected, although evidence in this regard is less consistent than in the case of vemurafenib for



(1) Tyrosine kinase inhibitor

FIGURE 2. A therapeutic algorithm for advanced melanoma. Tumor molecular genotyping determines the options in first-line treatment. Kinase inhibitors are the therapy of choice if a specific mutation is detected, although data are more consistent for vemurafenib in *BRAF*-mutant melanoma than for KIT inhibitors in *KIT*-mutant tumors. Ipilimumab, chemotherapy, or immunotherapy can be considered in nonmutated melanoma as first line or in mutated melanoma after progression with specific drugs. It is very rarely possible to use ipilimumab after progression on BRAF inhibitors; the use of these drugs in the first line cannot be excluded.

BRAF-mutant melanoma. Ongoing studies will determine which *KIT* mutations are most amenable to treatment with this kind of inhibitors.

Patients progressing on vemurafenib or a KIT inhibitor could receive ipilimumab or chemotherapy, depending on performance status and drug availability. Ipilimumab should also be considered as first-line therapy whenever a targetable mutation is not detected in the tumor. Immunotherapy with interleukin-2 or adoptive cell therapy should be restricted to selected patients in centers with experience in these treatment modalities.

The importance of clinical trials cannot be overemphasized. The possibility to refer patients for clinical investigation should be considered at any stage of the patient's evolution. The algorithm will be subjected to changes as new alternatives demonstrate efficacy for specific subgroups. For instance, MEK inhibitors or the

combination of BRAF and MEK inhibitors might be considered in first-line for *BRAF*-mutated melanomas. In the future, specific drugs could be developed to treat melanoma with other mutations, in which the detection of these mutations should become the standard of practice. Likewise, if markers predicting response to ipilimumab or anti-PD1 antibodies are finally found, they should be incorporated to the pathological workup.¹⁶

CONCLUSIONS

In 2012, therapy for advanced melanoma is determined by the mutational status of the tumor. Vemurafenib is the most active agent in *BRAF*-mutant melanoma, whereas a KIT inhibitor could be considered in tumors with a mutation in *KIT*. Ipilimumab should be

considered as first-line for patients with no target mutations or as second line in any patient. Chemotherapy may have a role as salvage therapy in second or third line.

REFERENCES

1. Balch CM, Soong SJ, Murad TM, et al. A multifactorial analysis of melanoma. IV. Prognostic factors in 200 melanoma patients with distant metastases (stage III). *J Clin Oncol.* 1983;1:126–134.
2. Manola J, Atkins M, Ibrahim J, et al. Prognostic factors in metastatic melanoma: a pooled analysis of Eastern Cooperative Oncology Group trials. *J Clin Oncol.* 2000;18:3782–3793.
3. Eigentler TK, Caroli UM, Radny P, et al. Palliative therapy of disseminated malignant melanoma: a systematic review of 41 randomised clinical trials. *Lancet Oncol.* 2003;4:748–759.
4. Patel PM, Suci S, Mortier L, et al. Extended schedule, escalated dose temozolomide versus dacarbazine in stage IV melanoma: final results of a randomised phase III study (EORTC 18032). *Eur J Cancer.* 2011;47:1476–1483.
5. Middleton MR, Grob JJ, Aaronson N, et al. Randomized phase III study of temozolomide versus dacarbazine in the treatment of patients with advanced metastatic malignant melanoma. *J Clin Oncol.* 2000;18:158–166.
6. Quereux G, Dreno B. Fotemustine for the treatment of melanoma. *Expert Opin Pharmacother.* 2011;12:2891–2904.
7. Atkins MB, Lotze MT, Dutcher JP, et al. High-dose recombinant interleukin 2 therapy for patients with metastatic melanoma: analysis of 270 patients treated between 1985 and 1993. *J Clin Oncol.* 1999;17:2105–2116.
8. Dudley ME, Wunderlich JR, Yang JC, et al. Adoptive cell transfer therapy following non-myeloablative but lymphodepleting chemotherapy for the treatment of patients with refractory metastatic melanoma. *J Clin Oncol.* 2005;23:2346–2357.
9. Rosenberg SA, Dudley ME. Adoptive cell therapy for the treatment of patients with metastatic melanoma. *Curr Opin Immunol.* 2009;21:233–240.
10. Robert C, Ghiringhelli F. What is the role of cytotoxic T lymphocyte-associated antigen 4 blockade in patients with metastatic melanoma? *Oncologist.* 2009;14:848–861.
11. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med.* 2010;363:711–723.
12. Chapman PB, Hauschild A, Robert C, et al. Phase III randomized, open-label, multicenter trial (BRIM3) comparing BRAF inhibitor vemurafenib with dacarbazine (DTIC) in patients with V600EBRAF-mutated melanoma. *J Clin Oncol.* 2011;29(Suppl):LBA4.
13. Shahabi V, Whitney G, Hamid O, et al. Assessment of association between BRAF-V600E mutation status in melanomas and clinical response to ipilimumab. *Cancer Immunol Immunother.* 2012;61:733–737.
14. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin Cancer Res.* 2009;15:7412–7420.
15. Brahmer JR, Horn L, Antonia S, et al. Clinical activity and safety of anti-PD1 (BMS-936558, MDX-1106) in patients with advanced non-small-cell lung cancer (NSCLC). *J Clin Oncol.* 2012;30(Suppl):7509.
16. Brahmer JR, Tykodi SS, Chow LQ, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med.* 2012;366:2455–2465.
17. Long GV, Menzies AM, Nagrial AM, et al. Prognostic and clinicopathologic associations of oncogenic BRAF in metastatic melanoma. *J Clin Oncol.* 2011;29:1239–1246.
18. Lovly CM, Dahlman KB, Fohn LE, et al. Routine multiplex mutational profiling of melanomas enables enrollment in genotype-driven therapeutic trials. *PLoS One.* 2012;7:e35309.
19. Meier F, Schitteck B, Busch S, et al. The RAS/RAF/MEK/ERK and PI3K/AKT signaling pathways present molecular targets for the effective treatment of advanced melanoma. *Front Biosci.* 2005;10:2986–3001.
20. Lee JT, Li L, Brafford PA, et al. PLX4032, a potent inhibitor of the B-Raf V600E oncogene, selectively inhibits V600E-positive melanomas. *Pigment Cell Melanoma Res.* 2010;23:820–827.
21. Yang H, Higgins B, Kolinsky K, et al. RG7204 (PLX4032), a selective BRAFV600E inhibitor, displays potent antitumor activity in preclinical melanoma models. *Cancer Res.* 2010;70:5518–5527.
22. Yancovitz M, Litterman A, Yoon J, et al. Intra- and inter-tumor heterogeneity of BRAF(V600E) mutations in primary and metastatic melanoma. *PLoS One.* 2012;7:e29336.
23. Sosman JA, Kim KB, Schuchter L, et al. Survival in BRAF V600-mutant advanced melanoma treated with vemurafenib. *N Engl J Med.* 2012;366:707–714.
24. Dummer R, Rinderknecht J, Goldinger SM. Ultraviolet A and photosensitivity during vemurafenib therapy. *N Engl J Med.* 2012;366:480–481.
25. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med.* 2011;364:2507–2516.
26. Dummer R, Rinderknecht J, Goldinger SM, et al. An open-label pilot study of vemurafenib in previously treated metastatic melanoma patients with brain metastases. *J Clin Oncol.* 2011;29(Suppl):8548.
27. Atkins MB. A randomized phase III trial of vemurafenib vs. ipilimumab in patients with advanced BRAFV 600 mutant melanoma. Available at: <http://ecog-acrin.org/trials/melanoma>. Accessed April 23, 2013.
28. Bristol-Myers Squibb. NCT01400451: a phase I/II trial of vemurafenib and ipilimumab in subjects with V600 BRAF mutation-positive metastatic melanoma. Available at: <http://clinicaltrials.gov/ct2/show/NCT01400451?term=NCT01400451&rank=1>. Accessed April 23, 2013.
29. Falchook GS, Long GV, Kurzrock R, et al. Dabrafenib in patients with melanoma, untreated brain metastases, and other solid tumours: a phase 1 dose-escalation trial. *Lancet.* 2012;379:1893–1901.
30. Hauschild A, Grob JJ, Demidov LV, et al. Phase III, randomized, open-label, multicenter trial (BREAK-3)

- comparing the BRAF kinase inhibitor dabrafenib (GSK2118436) with dacarbazine (DTIC) in patients with BRAFV600E-mutated melanoma. *J Clin Oncol*. 2012;30(Suppl):LBA8500.
31. Kirkwood JM, Long GV, Trefzer U, et al. BREAK-MB: a phase II study assessing overall intracranial response rate (OIRR) to dabrafenib (GSK2118436) in patients (pts) with BRAF V600E/k mutation-positive melanoma with brain metastases (mets). *J Clin Oncol*. 2012;30(Suppl):8501.
 32. Kaplan FM, Shao Y, Mayberry MM, et al. Hyperactivation of MEK-ERK1/2 signaling and resistance to apoptosis induced by the oncogenic B-RAF inhibitor, PLX4720, in mutant N-RAS melanoma cells. *Oncogene*. 2011;30:366–371.
 33. Ribas A, Flaherty KT. BRAF targeted therapy changes the treatment paradigm in melanoma. *Nat Rev Clin Oncol*. 2011;8:426–433.
 34. Infante JR, Falchook GS, Lawrence DP, et al. Phase I/II study to assess safety, pharmacokinetics, and efficacy of the oral MEK 1/2 inhibitor GSK1120212 (GSK212) dosed in combination with the oral BRAF inhibitor GSK2118436 (GSK436). *J Clin Oncol*. 2011;29(Suppl):CRA8503.
 35. Robert C, Flaherty KT, Hersey P, et al. METRIC phase III study: efficacy of trametinib (T), a potent and selective MEK inhibitor (MEKi), in progression-free survival (PFS) and overall survival (OS), compared with chemotherapy (C) in patients (pts) with BRAFV600E/K mutant advanced or metastatic melanoma (MM). *J Clin Oncol*. 2012;30(Suppl):LBA8509.
 36. Emery CM, Vijayendran KG, Zipser MC, et al. MEK1 mutations confer resistance to MEK and B-RAF inhibition. *Proc Natl Acad Sci U S A*. 2009;106:20411–20416.
 37. Weber JS, Flaherty KT, Infante JR, et al. Updated safety and efficacy results from a phase I/II study of the oral BRAF inhibitor dabrafenib (GSK2118436) combined with the oral MEK 1/2 inhibitor trametinib (GSK1120212) in patients with BRAFi-naïve metastatic melanoma. *J Clin Oncol*. 2012;30(Suppl):8510.
 38. Atefi M, von Euw E, Attar N, et al. Reversing melanoma cross-resistance to BRAF and MEK inhibitors by co-targeting the AKT/mTOR pathway. *PLoS One*. 2011;6:e28973.
 39. Greger JG, Eastman SD, Zhang V, et al. Combinations of BRAF, MEK, and PI3K/mTOR inhibitors overcome acquired resistance to the BRAF inhibitor GSK2118436 dabrafenib, mediated by NRAS or MEK mutations. *Mol Cancer Ther*. 2012;11:909–920.
 40. Nazarian R, Shi H, Wang Q, et al. Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature*. 2010;468:973–977.
 41. Poulikakos PI, Zhang C, Bollag G, et al. RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF. *Nature*. 2010;464:427–430.
 42. Wagle N, Emery C, Berger MF, et al. Dissecting therapeutic resistance to RAF inhibition in melanoma by tumor genomic profiling. *J Clin Oncol*. 2011;29:3085–3096.
 43. Schoenewolf NL, Bull C, Belloni B, et al. Sinonasal, genital and acrolentiginous melanomas show distinct characteristics of KIT expression and mutations. *Eur J Cancer*. 2012;48:1842–1852.
 44. Beadling C, Jacobson-Dunlop E, Hodi FS, et al. KIT gene mutations and copy number in melanoma subtypes. *Clin Cancer Res*. 2008;14:6821–6828.
 45. Guo J, Si L, Kong Y, et al. Phase II, open-label, single-arm trial of imatinib mesylate in patients with metastatic melanoma harboring c-Kit mutation or amplification. *J Clin Oncol*. 2011;29:2904–2909.
 46. Carvajal RD, Antonescu CR, Wolchok JD, et al. KIT as a therapeutic target in metastatic melanoma. *JAMA*. 2011;305:2327–2334.
 47. Ascierto PA, Berking C, Agarwala SS, et al. Efficacy and safety of oral MEK162 in patients with locally advanced and unresectable or metastatic cutaneous melanoma harboring BRAFV600 or NRAS mutations. *J Clin Oncol*. 2012;30(Suppl):8511.
 48. Dummer R, Guggenheim M, Arnold AW, et al. Updated Swiss guidelines for the treatment and follow-up of cutaneous melanoma. *Swiss Med Wkly*. 2011;141:w13320.
 49. Anderson S, Bloom KJ, Vallera DU, et al. Multisite analytic performance studies of a Real-time polymerase chain reaction assay for the detection of BRAF V600E mutations in formalin-fixed paraffin-embedded tissue specimens of malignant melanoma. *Arch Pathol Lab Med*. 2012;136:1385–1391.